arXiv:1101.0493v1 [math.AG] 3 Jan 2011

Monodromy of A-hypergeometric functions

Frits Beukers

December 20, 2010

Abstract

Using Mellin-Barnes integrals we give a method to compute a relevant subgroup of the monodromy group of an A-hypergeometric system of differential equations. Presumably this group is the full monodromy group of the system

1 Introduction

At the end of the 1980's Gel'fand, Kapranov and Zelevinsky, in [10], [11], [12], defined a general class of hypergeometric functions, encompassing the classical one-variable hypergeometric functions, the Appell and Lauricella functions and Horn's functions. They are called A-hypergeometric functions and they provide a beautiful and elegant basis of a theory of hypergeometric functions in several variables. For an introduction to the subject we refer the reader to [29], [5] or the book by Saito, Sturmfels and Takayama, [26]. We briefly recall the main facts. Let $A \subset \mathbb{Z}^r$ be a finite set such that

1. The $\mathbb{Z}$-span of A is $\mathbb{Z}^r$.
2. There exists a linear form h such that $h(\mathbf{a}) = 1$ for all $\mathbf{a} \in A$.

Let $\alpha = (\alpha_1, \ldots, \alpha_r) \in \mathbb{R}^r$. Denote $A = \{\mathbf{a}_1, \ldots, \mathbf{a}_N\}$ (with $N > r$). Writing the vectors $\mathbf{a}_i$ in column form we get the so-called A-matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1N} \\ a_{21} & a_{22} & \cdots & a_{2N} \\ \vdots & & & \vdots \\ a_{r1} & a_{r2} & \cdots & a_{rN} \end{pmatrix}$$

For $i = 1, 2, \ldots, r$ consider the first order differential operators

$$Z_i = a_{i1} v_1 \partial_1 + a_{i2} v_2 \partial_2 + \cdots + a_{iN} v_N \partial_N$$

where $\partial_j = \frac{\partial}{\partial v_j}$ for all j.

Let

$$L = \{(l_1, \dots, l_N) \in \mathbb{Z}^N \mid l_1\mathbf{a}_1 + l_2\mathbf{a}_2 + \cdots + l_N\mathbf{a}_N = \mathbf{0}\}$$

be the lattice of integer relations between the elements of A. For every $\mathbf{l} \in L$ we define the so-called box-operator

$$\Box_{\mathbf{l}} = \prod_{l_i>0} \partial_i^{l_i} - \prod_{l_i<0} \partial_i^{-l_i}$$

The system of differential equations

$$\begin{array}{rcll} (Z_i - \alpha_i)\Phi & = & 0 & (i = 1, \dots, r) \\ \Box_{\mathbf{l}}\Phi & = & 0 & \mathbf{l} \in L \end{array}$$

is known as the system of *A-hypergeometric differential equations* and we denote it by $H_A(\alpha)$. It turns out that the solution space of $H_A(\alpha)$ is finite dimensional with dimension equal to the volume of the convex hull $Q(A)$ of A. More precisely,

Theorem 1.1 (GKZ, Adolphson) *Suppose either one of the following conditions holds,*

1. *the toric ideal I_A in $\mathbb{C}[\partial_1, \dots, \partial_N]$ generated by the box operators has the Cohen-Macaulay property.*
2. *The system $H_A(\alpha)$ is non-resonant.*

Then the rank of $H_A(\alpha)$ is finite and equals the volume of the convex hull $Q(A)$ of the points of A. The volume is normalized so that a minimal $(r-1)$-simplex with integer vertices in $h(\mathbf{x}) = 1$ has volume 1.

Let $C(A)$ be the cone generated by the $\mathbb{R}_{\ge 0}$-linear combinations of $\mathbf{a}_1, \dots, \mathbf{a}_N$. We say that an A-hypergeometric system is *non-resonant* if the boundary of $C(A)$ has empty intersection with the shifted lattice $\alpha + \mathbb{Z}^r$. Theorem 1.1 is proven in [12], (corrected in [14]) and [1, Corollary 5.20].

Among the many papers written on A-hypergeometric equations there are virtually no papers dealing with the monodromy group of these systems. In the case of one-variable hypergeometric functions there is [8] which give an characterisation of monodromy groups as complex reflection groups. There is also a classical method to compute monodromy with respect to an explicit basis of functions using so-called Mellin-Barnes integrals, see F.C.Smith, [28]. A recent paper by Golyshev and Mellit, [15], deals with the same problem using Fourier-transforms of Γ-products. A recent paper by K.Mimachi [23] uses computation of twisted cycle intersection.

For the two-variable Appell system F_1 and Lauricella's F_D monodromy follows from the work by E.Picard [25], T.Terada [31] and Deligne-Mostow [9]. In T.Sasaki's paper [27] we find explicit monodromy generators for Appell F_1. They all use the fact that Lauricella functions of F_D-type can be written as one-dimensional twisted period integrals and monodromy is a representation of a braid group.

The two-variable Appell F_2 has been considered explicitly by M.Kato, [18]. The Appell F_3 system has the same A-set as Appell F_2, and therefore gives nothing new. Finally, the Appell system F_4 has been considered completely explicitely by K.Takano [30] and later Kaneko, [17]. In Haraoka, Ueno [16] we find some rigidity considerations on the monodromy of F_4.

Finally, the complete monodromy of the Aomoto-Gel'fand system $E(3,6)$ has been determined by K.Matsumoto, T.Sasaki, N.Takayama, M.Yoshida in [21] and further properties in [22]. See also M.Yoshida's book 'Hypergeometric Functions, my Love', [32].

It is the purpose of the present paper to compute monodromy groups of A-hypergeometric systems by methods which are combinatorial in nature. We do this by starting with local monodromy groups which arise from series expansions of solutions of $H_A(\alpha)$. It is well known that such local expansions correspond one-to-one with regular triangulations of A. This is a discovery by Gel'fand, Kapranov and Zelevinsky that we shall explain in Section 5. The local monodromy groups have to be glued together to build a global monodromy group. This glue is provided by multidimensional Mellin-Barnes integrals as defined in Section 2. If the Mellin-Barnes integrals provide a basis of local solutions (Assumption 4.5), the construction of the global group generated by the local contributions is completely combinatorial. We call this group $lMon$ and denote the global monodromy group by Mon. We now make the following hypothesis

Hypothesis 1.2

$$lMon = Mon$$

Assuming this hypothesis we can compute monodromy groups in a combinatorial way. As in [8] this description is up to common conjugation of the monodromy matrices. If one would like to calculate monodromy matrices with respect to explicit bases of solutions one would have to find an explicit calculation of Mellin-Barnes integral as a linear combination of power series solutions. This is a tedious task which we like to carry out in a forthcoming paper. We remark that such a calculation has been carried out in the so-called confluent case (i.e. A does not lie in translated hyperplane) by O.N.Zhdanov and A.K.Tsikh, [33]. In her PhD-thesis from 2009, Lisa Nilsson [24] introduced (non-confluent) A-hypergeometric functions in terms of Mellin-

Barnes integrals and initiated their study. It is these integrals that we shall use.

In the remainder of this paper we assume that *the shifted lattice* $\alpha + \mathbb{Z}^r$ *has empty intersection with any hyperplane spanned by* $r-1$ *independent elements of* A. In that case we say that the system is *totally non-resonant.* Note that this is stronger than just non-resonance where only the faces of the cone spanned by the elements of A are involved. Non-resonance (and a fortiori total non-resonance) ensures that our system is irreducible, see for example [13, Thm 2.11] or [6] for a slightly more elementary proof. Non-resonance also implies that A-hypergeometric systems whose parameter vectors are the same modulo $\mathbb{Z}^r$ have isomorphic monodromy, see [6, Thm 2.1], which is actually a Theorem due to B.Dwork. Total non-resonance implies T-nonresonance for every triangulation T in the terminology of Gel'fand, Kapranov and Zelevinsky. In particular this implies that the local solution expansions will not contain logarithms. So local monodromy representations act by characters. We prefer to leave the case of logarithmic local solutions for a later occasion.

2 Mellin-Barnes integrals

Let notation be as usual with the additional notation $d = N - r$. This number will be the number of essential variables in the hypergeometric system (e.g. $d = 2$ in the Appell systems), the other $N - d$ variables exist for homogeneity purposes in the A-hypergeometruic setup. Define $\gamma_1, \ldots, \gamma_N$ such that $\gamma_1 \mathbf{a}_1 + \cdots + \gamma_N \mathbf{a}_N = \alpha$. We choose parameters $\mathbf{s} = (s_1, \ldots, s_d)$ on $L(\mathbb{R}) = L \otimes \mathbb{R}$ as follows. Choose a $\mathbb{Z}$-basis for the lattice L and write the basis elements as rows of a $d \times N$ matrix B, which we call the B-matrix. The matrix B has the property that it has maximal rank d and $A.B$ is the zero matrix. The columns of B can be considered as coordinates of the linear forms l_i on L or $L \otimes \mathbb{R}$. Denote these columns by $\mathbf{b}_j$. Then the space $L(\mathbb{R}) \subset \mathbb{R}^N$ is parametrized by the N-tuple $(\mathbf{b}_1 \cdot \mathbf{s}, \ldots, \mathbf{b}_N \cdot \mathbf{s})$ with $\mathbf{s} = (s_1, \ldots, s_d) \in \mathbb{R}^d$ as parameter vector.

We now complexify the parameters s_i and consider the integral

$$M(v_1, \ldots, v_N) = \int_{\sqrt{-1}\ \mathbb{R}^d} \prod_{i=1}^{N} \Gamma(-\gamma_i - \mathbf{b}_i \cdot \mathbf{s}) v_i^{\gamma_i + \mathbf{b}_i \cdot \mathbf{s}} d\mathbf{s} \qquad (MB)$$

where $d\mathbf{s} = ds_1 \wedge \cdots \wedge ds_d$ and the integration takes place over $\mathrm{Re}(s_i) = 0$ for $i = 1, \ldots, d$. This is an example of a so-called *Mellin-Barnes integral.* It will be crucial in the determination of the monodromy of A-hypergeometric systems.

3 Differential equations

We prove the following Theorem,

Theorem 3.1 *Assume that* $\gamma_i < 0$ *for* $i = 1, 2, \ldots, N$. *Then the Mellin-Barnes integral* $M(v_1, \ldots, v_N)$ *satisfies the set of A-hypergeometric equations* $H_A(\alpha)$.

This will be done under the assumption that the Mellin-Barnes integral converges absolutely. We come to the matter of convergence in the next section.

Proof. The Mellin-Barnes integral clearly has the property

$$M(\mathbf{t}^{\mathbf{a}_1} v_1, \ldots, \mathbf{t}^{\mathbf{a}_N} v_N) = \mathbf{t}^{\alpha} M(v_1, \ldots, v_N)$$

for all $\mathbf{t} \in (\mathbb{C}^*)^N$. So $M(\mathbf{v})$ satisfies the hypergeometric homogeneity equations.

Now let $\lambda \in L$ and put $\lambda = \lambda_+ - \lambda_-$. Define $|\lambda| = \sum_{i=1}^N |\lambda_i|$. Then

$$\begin{aligned}\Box_\lambda M(v_1, \ldots, v_N) &= (-1)^{|\lambda|/2} \int_{\sqrt{-1}\ \mathbb{R}^d} \prod_{i=1}^N \Gamma(-\gamma_i - \mathbf{b}_i \cdot \mathbf{s} + \lambda_{+,i}) v_i^{\gamma_i + \mathbf{b}_i \cdot \mathbf{s} - \lambda_{+,i}} d\mathbf{s} \\ &- (-1)^{|\lambda|/2} \int_{\sqrt{-1}\ \mathbb{R}^d} \prod_{i=1}^N \Gamma(-\gamma_i - \mathbf{b}_i \cdot \mathbf{s} + \lambda_{-,i}) v_i^{\gamma_i + \mathbf{b}_i \cdot \mathbf{s} - \lambda_{-,i}} d\mathbf{s}.\end{aligned}$$

Choose $\mathbf{s}_\lambda$ such that $\mathbf{b}_i \cdot \mathbf{s}_\lambda = \lambda_i$ for $i = 1, \ldots, N$. Then the second integral is actually integration over $\mathbf{s}_\lambda + \sqrt{-1}\ \mathbb{R}^d$ of the integrand of the first integral.

Because of the assumption $\gamma_i < 0$ we see that $-\gamma_i - t\mathbf{b}_i \cdot \mathbf{s}_\lambda + \lambda_{+,i} > 0$ for all $t \in [0, 1]$ and $i = 1, \ldots, N$. Hence the $d+1$-dimensional domain $\{t\mathbf{s}_\lambda + i\mathbb{R}^d | 0 \le t \le 1\}$ does not contain any poles of the integrand and a homotopy argument gives that the two integrals are equal and cancel. $\Box$

Not all systems $H_A(\alpha)$ allow a choice of negative γ_i. However, under the assumption of non-resonance it is known that two hypergeometric systems $H_A(\alpha)$ and $H_A(\alpha')$ have the same monodromy if $\alpha - \alpha' \in \mathbb{Z}^r$ (see [6, Theorem 2.1]). We call such systems contiguous. Thus we can always replace an irreducible A-hypergeometric system by a contiguous one which does allow a choice of $\gamma_i < 0$ for all i. From now on we make this assumption, i.e all our Mellin-Barnes integrals are solution of an A-hypergeometric system. Of course there is also the question whether or not $M(v_1, \ldots, v_N)$ is a trivial function. By Proposition 4.5 we will find that it is non-trivial.

4 Convergence

We find from [2] the following estimate. Suppose $s = a + bi$ with $a_1 < a < a_2$ and $|b| \to \infty$. Then

$$|\Gamma(a+bi)| = \sqrt{2\pi}|b|^{a-1/2}e^{-\pi|b|/2}[1+O(1/|b|)].$$

Notice also that for any $v \in \mathbb{C}^*$ we have $|v^{a+bi}| = |v|^a e^{-b\arg(v)}|$. Write $s_j = i\tau_j$ for $j = 1, \ldots, N-r$. Let us denote $\theta_j = \arg(v_j)$ and $l_j(\tau) = l_j(\tau_1, \ldots, \tau_d)$. The integrand in the Mellin-Barnes integral can now be estimated by

$$\left|\prod_{i=1}^{N} \Gamma(-\gamma_i - \mathbf{b}_i \cdot \mathbf{s}))v_i^{\gamma_i + \mathbf{b}_i \cdot \mathbf{s}}\right| \le c_1 \max_j |\tau_j|^{c_2} \exp\left(-\sum_{j=1}^{N}(\pi|\mathbf{b}_j \cdot \tau|/2 - \mathbf{b}_j \cdot \tau)\theta_j\right)$$

where c_1, c_2 are positive numbers depending only on γ_j, v_j. In order to ensure convergence of the integral we must have that $\sum_{j=1}^{N}(\pi|\mathbf{b}_j \cdot \tau|/2 + \mathbf{b}_j \cdot \tau))\theta_j > 0$ for every non-zero $\tau \in \mathbb{R}^d$. We apply the following Lemma.

Lemma 4.1 *Given N vectors $\mathbf{p}_1, \ldots, \mathbf{p}_N$ in $\mathbb{R}^d$. Suppose they have rank d. Let $\mathbf{q} \in \mathbb{R}^d$. Then the following statements are equivalent,*

i) For all non-zero $\mathbf{x} \in \mathbb{R}^d$:

$$|\mathbf{q} \cdot \mathbf{x}| < \sum_{i=1}^{N} |\mathbf{p}_i \cdot \mathbf{x}|$$

ii) There exist $\mu_1, \ldots, \mu_N$ with $-1 < \mu_i < 1$ such that

$$\mathbf{q} = \mu_1 \mathbf{p}_1 + \cdots + \mu_N \mathbf{p}_N.$$

Proof. First suppose that $\mathbf{q} = \mu_1 \mathbf{p}_1 + \cdots + \mu_N \mathbf{p}_N$. Then, for all non-zero $\mathbf{x} \in \mathbb{R}^d$,

$$\begin{aligned} |\mathbf{q} \cdot \mathbf{x}| &= |\sum_{i=1}^{N} \mu_i \mathbf{p}_i \cdot \mathbf{x}| \\ &\le \sum_{i=1}^{N} |\mu_i||\mathbf{p}_i \cdot \mathbf{x}| < \sum_{i=1}^{N} |\mathbf{p}_i \cdot \mathbf{x}| \end{aligned}$$

To show the converse statement consider the set

$$V = \left\{ \sum_{i=1}^{N} \mu_i \mathbf{p}_i \;\middle|\; -1 < \mu_i < 1 \right\}.$$

This is a convex set. Suppose $\mathbf{q} \notin V$. Then there exists a linear form h such that $h(\mathbf{q}) > h(\mathbf{p})$ for all $\mathbf{p} \in V$. In other words, there exists $\mathbf{x} \in \mathbb{R}^d$ such that $\mathbf{q} \cdot \mathbf{x} > \sum_{i=1}^N \lambda_i \mathbf{p}_i \cdot \mathbf{x}$ for all $-1 < \lambda_i < 1$. In particular,

$$|\mathbf{q} \cdot \mathbf{x}| \geq \sum_{i=1}^{N} |\mathbf{p}_i \cdot \mathbf{x}|$$

contradicting our assumption. Hence $\mathbf{q} \in V$.

□

Application of Lemma 4.1 to $\mathbf{q} = \sum_{j=1}^N \theta_j \mathbf{b}_j$ and $\mathbf{p}_j = \pi \mathbf{b}_j/2$ yields the following criterion.

Corollary 4.2 *Let notations be as above. Then the Mellin-Barnes integral converges absolutely if there exist* $\mu_i \in (-1,1)$ *such that*

$$\sum_{i=1}^{N} \frac{\theta_i}{2\pi} \mathbf{b}_i = \frac{1}{4} \sum_{i=1}^{N} \mu_i \mathbf{b}_i.$$

Let us define $Z_B = \{\frac{1}{4}\sum_{i=1}^N \mu_i \mathbf{b}_i | \mu_i \in (-1,1)\}$. This is a so-called zonotope in d-dimensional space. The convergence condition for the Mellin-Barnes integral now reads $\sum_{i=1}^N \frac{\theta_i}{2\pi}\mathbf{b}_i \in Z_B$.

As an example let us take the system Horn G_3. The set A is given by the columns of

$$A = \begin{pmatrix} 2 & 1 & 0 & -1 \\ -1 & 0 & 1 & 2 \end{pmatrix}.$$

The basisvectors $(1,-2,1,0),(0,1,-2,1)$ of L form the rows of a B-matrix

$$B = \begin{pmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \end{pmatrix}.$$

The parameter vector corresponding to G_3 is $(-a,-b)$. When $a, b > 0$ it is easy to see that there exist negative $\gamma_1, \ldots, \gamma_4$ such that $A(\gamma_1 \ldots \gamma_4)^t = (-a,-b)^t$. Let us assume we are in this situation. Thus the corresponding Mellin-Barnes integral is indeed a solution of the G_3-system. The zonotope Z_B can now be pictured as



and the convergence condition reads

$$\frac{1}{2\pi}(\theta_1 - 2\theta_2 + \theta_3, \theta_2 - 2\theta_3 + \theta_4) \in Z_B.$$

Note that the point $(-0.4, -0.4)$ as well as $(-0.4+1, -0.4)$ and $(-0.4, -0.4+1)$ are contained in Z_B. They correspond to the arguments $(\theta_1, \theta_2, \theta_3, \theta_4) = 2\pi(-0.4, 0, 0, -0.4)$, $2\pi(0.6, 0, 0, -0.4)$ and $2\pi(-0.4, 0, 0, 0.6)$. These argument choices represent the same point in v_1, v_2, v_3, v_4-space. Hence we have three Mellin-Barnes solutions of the system G_3 around one point. According to Proposition 4.3 these integrals are linearly independent. and hence form a basis of local solutions of the G_3-system. We say that we have a *Mellin-Barnes basis* of solutions.

Proposition 4.3 *Let* $v = (v_1, \dots, v_N) \in (\mathbb{C}^*)^N$ *and let* Θ *be a finite set of* N*-tuples* $\theta = (\theta_1, \dots, \theta_N)$ *such that* $v_i = |v_i| \exp(\theta_i)$ *and the sums* $\frac{1}{2\pi}\sum_{i=1}^{d} \theta_i \mathbf{b}_i$ *are distinct elements of* Z_B*. To each* $\theta \in \Theta$ *denote the corresponding determination of the Mellin-Barnes integral in the neighbourhood of* v *by* M_θ*.*

Then the functions M_θ *are linearly independent over* $\mathbb{C}$*.*

The proof of this Lemma depends on a d-dimensional version or if one wants, repeated application, of the following Theorem.

Theorem 4.4 (Mellin inversion theorem) *Let* $\phi(z)$ *be function on* $\mathbb{C}$ *satisfying the follwoing properties*

(a) ϕ *is analytic in a vertical strip of the form* $\alpha < x = \mathrm{Re}(z) < \beta$ *where* $\alpha, \beta \in \mathbb{R}$*.*

(b) $\int_{\infty}^{\infty} |\phi(x+iy)| dy$ *converges for all* $x \in (\alpha, \beta)$*.*

(c) $\phi(z) \to 0$ *uniformaly as* $|y| \to \infty$ *in* $\alpha + \epsilon < x < \beta - \epsilon$ *for all* $\epsilon > 0$*.*

Denote for all $\theta > 0$,

$$f(\theta) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \theta^{-z} \phi(z) dz.$$

Then,

$$\phi(z) = \int_0^\infty \theta^{z-1} f(\theta) d\theta.$$

For a proof of this theorem see [20, Appendix 4, p341-342]. A treatment of the multidimensional case can also be found for example in [3].

Proof of Proposition 4.3. Suppose we have a non-trivial relation $\sum_{\theta\in\Theta} \lambda_\theta M_\theta = 0$. Let us use the notation $(\theta \cdot \mathbf{b})(\mathbf{s}) = \theta_1 \mathbf{b}_1 \cdot \mathbf{s} + \cdots + \theta_N \mathbf{b}_N \cdot \mathbf{s}$. The relation can be written as

$$0 = \int_{\mathbf{s}\in\sqrt{-1}\mathbb{R}^d} \left(\sum_{\theta\in\Theta} \lambda_\theta e^{(\theta\cdot\mathbf{b})(\mathbf{s})} \right) |v_1|^{\mathbf{b}_1\cdot\mathbf{s}} \cdots |v_N|^{\mathbf{b}_N\cdot\mathbf{s}} \prod_{i=1}^N \Gamma(-\gamma_i - \mathbf{b}_i \cdot \mathbf{s}) d\mathbf{s}$$

Let us now write $x_j = |v_1|^{b_{1j}} \cdots |v_N|^{b_{Nj}}$ where b_{ij} are the entries of the B-matrix. Then

$$|v_1|^{\mathbf{b}_1\cdot\mathbf{s}} \cdots |v_N|^{\mathbf{b}_N\cdot\mathbf{s}} = x_1^{s_1} \cdots x_d^{s_d}$$

By repeated use of the Mellin inversion Theorem 4.4 we conclude that the vanishing of the integral implies the identical vanishing of

$$\sum_{\theta\in\Theta} \lambda_\theta e^{(\theta\cdot\mathbf{b})(\mathbf{s})}$$

Since the exponentials are all distinct functions of $s_1, \ldots, s_d$ this implies that $\lambda_\theta = 0$ for all $\theta \in \Theta$.

□

An alternative proof of Proposition 4.3 can be found in [24, Lemma 5.5].

Since we assume that $\gamma_i < 0$ for all i, all Mellin-Barnes integrals are solutions of the corresponding A-hypergeometric system. It would be very convenient if such a basis of solutions given by Mellin-Barnes integrals would always exist. It turns out that with the exception of Appell F_4 all $d = 2$ systems Appell F_1, F_2, F_3 and Horn $G_1, G_2, G_3, H_1, \ldots, H_7$ this is the case. A theoretical framework for a result like this may be provided in the PhD-thesis of Lisa Nilsson [24] suggesting that there does indeed exist such a basis if the complement of the so-called coamoeba of the A-resultant is non-empty. In [24] this is elaborated for the case $d = 2$.

Let us now make the following assumption on our system $H_A(\alpha)$.

Assumption 4.5 *There exists a point $v \in (\mathbb{C}^*)^N$ around which we can find a Mellin-Barnes basis of solutions.*

For the practical determination of a Mellin-Barnes basis we use the following Proposition.

Proposition 4.6 *Let $H_A(\alpha)$ be a totally non-resonant system of rank D. The system allows a Mellin-Barnes basis of solutions if and only if the zonotope Z_B contains D distinct points $\tau_1, \ldots, \tau_D$ whose coordinates differ by integers.*

Proof. From the discussion above it follows that the existence of a Mellin-Barnes basis corresponds to the choice of D N-tuples $(\theta_1, \ldots, \theta_N)$ representing argument choices of a given point. Hence the differences between these N-tuples have coordinates which are integer multiples of 2π. The sums $\frac{1}{2\pi}\sum_{i=1}^N \theta_i \mathbf{b}_i$ are distinct, hence the D Mellin-Barnes basis elements correspond to D points $\tau_i \in Z_B$ whose coordinates differ by integers.

Suppose conversely we have D points $\tau_i \in Z_B$ whose coordinates differ by integers. Since the $\mathbb{Z}$-span of the columns $\mathbf{b}_i$ is $\mathbb{Z}^d$ we can find for every i integers $n_{i1}, \ldots, n_{iN}$ such that $\tau_i - \tau_1 = n_{i1}\mathbf{b}_1 + \cdots + n_{iN}\mathbf{b}_N$. So if $(\theta_1, \ldots, \theta_N)$ is an argument choice for τ_1, then the N-tuples $(\theta_1 + 2\pi n_{i1}, \ldots, \theta_N + 2\pi n_{iN})$ represent argument choices for τ_i with $i = 1, 2, \ldots, N$.

□

In practice we proceed as follows. We assume that there exists $I \subset \{1, \ldots, N\}$ of cardinality d such that $\det(\mathbf{b}_i)_{i\in I} = \pm 1$. This happens in almost all cases of interest. Then to any $\tau \in Z_B$ we can choose $(\theta_1, \ldots, \theta_N)$ with $\theta_i = 0$ for all $i \notin I$ such that $2\pi\tau = \sum_{i=1}^N \theta_i \mathbf{b}_i$. From the determinant condition it follows that integer coordinate differences for the points τ correspond to integer coordinate differences in the $\theta_i/2\pi$.

5 Power series solutions

Consider the system $H_A(\alpha)$ and a formal solution

$$\Phi_\gamma = \sum_{\mathbf{l}\in L} \frac{v_1^{l_1+\gamma_1} \cdots v_N^{l_N+\gamma_N}}{\Gamma(l_1+\gamma_1+1)\cdots\Gamma(l_N+\gamma_N+1)}$$

where γ is chosen such that $\alpha = \gamma_1\mathbf{a}_1 + \cdots + \gamma_N\mathbf{a}_N$. This series expansion is a Laurent series multiplied by generally non-integral powers of the variables v_i. We call such a series a twisted Laurent series. As is well-known we have the freedom of choice in γ. We shall use this freedom in the following way where, again, we denote the columns of the B-matrix by $\mathbf{b}_i$. Choose a subset $I \subset \{1, 2, \ldots, N\}$ with $|I| = d = N - r$

such that $\mathbf{b}_i$ with $i \in I$ are linearly independent. It is known that $|\det(\mathbf{b}_i)_{i\in I}| = |\det(\mathbf{a}_j)_{j\notin I}|$ and we denote this quantity by Δ_I. Choose γ such that $\gamma_i \in \mathbb{Z}$ for all $i \in I$. There are precisely Δ_I such choices for γ which are distinct modulo L. The series Φ now reads

$$\Phi_\gamma = \sum_{\mathbf{l}\in L} \prod_{i\in I} \frac{v_i^{l_i+\gamma_i}}{\Gamma(l_i+\gamma_i+1)} \times \prod_{j\notin I} \frac{v_j^{l_j+\gamma_i}}{\Gamma(l_j+\gamma_j+1)}$$

Since $\gamma_i \in \mathbb{Z}$ for all $i \in I$ this summation extends over all $\mathbf{l}$ with $\gamma_i + l_i \geq 0$ for all $i \in I$. The other terms vanish because $1/\Gamma(\gamma_i + l_i + 1) = 0$ whenever $\gamma_i + l_i$ is a negative integer for some $i \in I$. Hence Φ_γ is now a twisted power series. Let us fix such a choice of γ. It is not hard to see that for γ-values which are distinct modulo L, the corresponding twisted power series are linear independent over $\mathbb{C}$.

Let now $\rho_1, \ldots, \rho_N$ be any N-tuple with the property that $\rho_1 l_1 + \cdots + \rho_N l_N > 0$ for any non-zero $\mathbf{l} \in L$ with $l_i \geq 0$ for all $i \in I$. Then, according to Theorem [5, Proposition 16.2] or [29, Section 3.3,3.4], the series Φ_γ converges for all $v_1, \ldots, v_N$ with $\forall i : |v_i| = t^{\rho_i}$ and $t \in \mathbb{R}_{>0}$ sufficiently small. We call such an N-tuple $\rho_1, \ldots, \rho_N$ a *convergence direction* of Φ_γ.

There is one important assumption we need in order to make this approach work. Namely the garantee that none of the arguments $\gamma_j + l_j$ is a negative integer when $j \notin I$. Otherwise we might even end up with a trivial series solution. Notice that

$$\alpha = \sum_{j=1}^{N} (\gamma_j + l_j)\mathbf{a}_i \equiv \sum_{j\notin I} \gamma_j \mathbf{a}_j (\text{mod } \mathbb{Z}^r)$$

So if $\gamma_j \in \mathbb{Z}$ for some $j \notin I$, the point α lies modulo $\mathbb{Z}^r$ in a space spanned by the $d-1$ remaining vectors $\mathbf{a}_i$. Under the assumption of total non-resonance this situation cannot occur.

So we recall our assumption that $H_A(\alpha)$ is totally non-resonant. We denote the set of all sets I such that $\Delta_I = |\det(\mathbf{b}_i)_{i\in I}| \neq 0$ by $\mathcal{I}$. We agree that we allow $s = \Delta_I$ copies of I in this list. To each $I \in \mathcal{I}$ there corresponds a choice of γ and we see to it that all these choices are distinct modulo L. So to an index set I which occurs s times there correspond s choices of γ that are distinct modulo L. The corresponding powerseries solutions are denoted by Φ_I.

Choose $I \in \mathcal{I}$ and a convergence direction $\rho = (\rho_1, \ldots, \rho_N)$ such that $\rho_1 l_1 + \cdots + \rho_N l_N > 0$ for all non-zero $\mathbf{l} \in L$ with $\forall i \in I : l_i \geq 0$. We can consider ρ as a linear form on $L \otimes \mathbb{R}$ and hence as an element in the span of the columns $\mathbf{b}_i$ of the B-matrix. From the positivity condition we infer that ρ must be in the strictly positive cone spanned by the forms given by $\mathbf{b}_i, i \in I$. Let us denote

$$\mathbf{b}_I = \left\{ \sum_{i \in I} \lambda_i \mathbf{b}_i \,\middle|\, \lambda_i > 0 \right\}.$$

Then the convergence directions with respect to I are given by the elements of $\mathbf{b}_I$. Conversely, fix an element ρ in the span of all $\mathbf{b}_i$ which does not lie on the boundary of any $\mathbf{b}_I$. Define $\mathcal{I}(\rho) = \{I|\rho \in \mathbf{b}_I\}$. Then, by the theory of Gel'fand, Kapranov and Zelevinsky the powerseries Φ_I with $I \in \mathcal{I}(\rho)$ form a basis of solutions with a common open region of convergence. We call such a set a basis of local solutions of $H_A(\alpha)$. It also follows from the theory that the sets $\mathcal{I}(\rho)$ are in one-to-one correspondence with the regular triangulations of the set A. This correspondence is given by $\mathcal{I}(\rho) \mapsto \{I^c|I \in \mathcal{I}(\rho)\}$.

The intersections of $\mathbf{b}_I$ define a subdivision of $\mathbb{R}^d$ into open convex polyhedral cones whose closure of the union is $\mathbb{R}^d$. This is a polyhedral fan which is called the *secondary fan*. The open cones in the secondary fan are in one-to-one correspondence with the bases of local series solutions.

Suppose we have a basis of local series solutions and suppose they converge in an open neighbourhood of a set defined by $|v_1| = r_1, \ldots, |v_N| = r_N$ with $r_i > 0$ for all i. Let $\gamma^{(1)}, \ldots, \gamma^{(D)}$ be the choices of γ used in the construction of the local series solutions $\Phi_1, \ldots, \Phi_D$. Given any N-tuple of integers $n_1, \ldots, n_N$ we define the closed path $c(n_1, \ldots, n_N)$ by

$$(r_1 e^{2\pi\sqrt{-1}\ n_1 t}, \ldots, r_N e^{2\pi\sqrt{-1}\ n_N t}), \quad t \in [0,1].$$

Taking $(r_1, \ldots, r_N)$ as a base point, the series Φ_j changes into

$$\exp(2\pi\sqrt{-1}\ (n_1\gamma_1^{(j)} + \cdots + n_N\gamma_N^{(j)})) = \exp(2\pi\sqrt{-1}\ \mathbf{n} \cdot \gamma^{(j)})$$

times Φ_j after analytic continuation along $c(n_1, \ldots, n_N)$. The group of these substitutions is called a *local monodromy group* with respect to $\Phi_1, \ldots, \Phi_D$.

6 Monodromy computation

Suppose our system $H_A(\alpha)$ allows a Mellin-Barnes basis of solutions (Assumption 4.5). Denote the basis elements by $M_1, \ldots, M_D$ and the corresponding points in Z_B by $\tau_1, \ldots, \tau_D$. It is the goal of this section to compute the local monodromy groups with respect to the basis $M_1, \ldots, M_D$. To this end we shall determine the transition matrices from $M_1, \ldots, M_D$ to each of the bases of local series expansions $\Phi_1, \ldots, \Phi_D$.

To each point τ_i there corresponds a (not necessarily unique) choice of arguments $\Theta_i = (\mathrm{Arg}(v_1), \dots, \mathrm{Arg}(v_N))$ for $i = 1, \dots, D$. We assume that the arguments are chosen such that the differences $\Theta_i - \Theta_1$ have all their components equal to integer multiples of 2π (see the comments following Proposition 4.6). Let $v \in (\mathbb{C}^*)^N$ be a point where we have a Mellin-Barnes basis of solutions. The Mellin-Barnes integral corresponding to the argument vector Θ_i is denoted by M_i.

For any N-tuple of integers $\mathbf{n} = (n_1, \dots, n_N)$ we consider the loop

$$c = c(\mathbf{n}) = c(n_1, \dots, n_N) : \ (e^{2\pi\sqrt{-1}n_1 t})v_1, \dots, e^{2\pi\sqrt{-1}n_N t}), \ t \in [0, 1].$$

Note that after analytic continuation of M_1 along the path $c((\Theta_j - \Theta_1)/2\pi)$ we end up with the Mellin-Barnes solution M_j for every j. We denote this path by c_j.

Now let ρ be a convergence direction and consider the corresponding basis of local powerseries solutions. Suppose that the realm of convergence of the local series expansions contains the points with $|v_i| = r_i$, for all i. Choose a path from v to a point on $|v_i| = r_i$ by keeping the argument values fixed. Then follow the loop c_j and finally return to v. We perform analytic continuation of all M_i along this composite path which we call c_j again. Let us denote a basis of local series expansions by $f_1, \dots, f_D$ and the corresponding choices of γ by $\gamma^{(1)}, \dots, \gamma^{(D)}$. Then there exist scalars μ_i such that

$$M_1 = \mu_1 f_1 + \cdots + \mu_D f_D.$$

After continuation along c_j the integral M_1 changes into M_j for every j. Under the paths c_j the local expansions f_i are multiplied by scalars. The space spanned by $M_1, \dots, M_D$ is D-dimensional. The space spanned by the images of $\mu_1 f_1 + \cdots + \mu_D f_D$ under $c_j, j = 1, 2, \dots, D$ is at most equal to the number of non-zero μ_i. Hence we conclude that $\mu_i \neq 0$ for all i. Let us renormalise the f_i such that

$$M_1 = f_1 + \cdots + f_D.$$

Then after continuation along c_j we get

$$M_j = e^{\sqrt{-1}\ (\Theta_j - \Theta_1)\cdot\gamma^{(1)}} f_1 + \cdots + e^{\sqrt{-1}\ (\Theta_j - \Theta_1)\cdot\gamma^{(D)}} f_D$$

for every j. Define

$$X_\rho = \begin{pmatrix} 1 & \cdots & 1 \\ e^{\sqrt{-1}\ (\Theta_2 - \Theta_1)\cdot\gamma^{(1)}} & \cdots & e^{\sqrt{-1}\ (\Theta_2 - \Theta_1)\cdot\gamma^{(D)}} \\ \vdots & & \vdots \\ e^{\sqrt{-1}\ (\Theta_D - \Theta_1)\cdot\gamma^{(1)}} & \cdots & e^{\sqrt{-1}\ (\Theta_D - \Theta_1)\cdot\gamma^{(D)}} \end{pmatrix}.$$

Then

$$\begin{pmatrix} M_1 \\ \vdots \\ M_D \end{pmatrix} = X_\rho \begin{pmatrix} f_1 \\ \vdots \\ f_D \end{pmatrix}$$

hence X_ρ is the desired transition matrix. Let χ be a diagonal $D \times D$-matrix which is in the local monodromy representation of $f_1, \ldots, f_D$. With respect to $M_1, \ldots, M_D$ this monodromy element has matrix $X_\rho \chi X_\rho^{-1}$. Thus we see that all local monodromies can be written with respect to a fixed Mellin-Barnes basis.

7 Hermitian forms

Theorem 7.1 *Let $H_A(\alpha)$ be a non-resonant A-hypergeometric system with $\alpha \in \mathbb{R}^r$. Let V be a vectorspace of local powerseries solutions. Then there exists a non-trivial unique (up to scalars) Hermitean form on V which is invariant with respect to the monodromy group of the system.*

Proof. In [7] we find that the solution space can be given by integrals over suitable cycles of integration (generalised Pochhammer cycles) or over twisted (or loaded) cycles in the sense of Kita and Yoshida [19]. The Hermitean form arises from the intersection form on these cycles. □

Suppose that $\alpha \in \mathbb{R}^r$ and that $H_A(\alpha)$ is totally non-resonant. Suppose also that we have a Mellin-Barnes basis of solutions. It is the purpose of this section to find explicit expressions for this Hermitean form with respect to the Mellin-Barnes basis.

Let us recall the notations $\mathbf{b}_I, \mathcal{I}$ from Section 5. To each $I \in \mathcal{I}$ there is associated a choice of γ which we denote γ^I. Let $\Theta_1, \ldots, \Theta_D \in \mathbb{R}^N$ be the argument vectors corresponding to the Mellin-Barnes basis elements $M_1, \ldots, M_D$. To each $I \in \mathcal{I}$ we associate the vector

$$X_I = (1, \exp(\sqrt{-1}\,(\Theta_2 - \Theta_1)\cdot\gamma^I), \ldots, \exp(\sqrt{-1}\,(\Theta_D - \Theta_1)\cdot\gamma^I)^t \in \mathbb{C}^D. \tag{1}$$

Vectors of this form are the column vectors of the transition matrices X_ρ from the previous section.

To every monodromy element we associate $g \in GL(n, \mathbb{C})$ such that the monodromy substition has the form $(M_1, \ldots, M_D)^t \to g(M_1, \ldots, M_D)^t$. By Theorem 7.1 there exists a Hermitean $n \times n$-matrix H (i.e. $\overline{H}^t = H$) such that $\overline{g}^t H g = H$ for all monodromy matrices g.

Proposition 7.2 *Let notations be as above. Let $I, J \in \mathcal{I}$ and suppose that $\mathbf{b}_I$ and $\mathbf{b}_J$ have a point in common. Then X_I, X_J are orthogonal with respect to H, i.e $\overline{X}_J^t H X_I = 0$.*

Proof. Given a basis of local solutions we have seen in Section 6 that the local monodromies with respect to the Mellin-Barnes basis have the form $g = X_\rho \chi X_\rho^{-1}$ where χ is a diagonal matrix representation with entries $\mathbf{n} \in \mathbb{Z}^N \to (\exp(2\pi\sqrt{-1}\,\mathbf{n}\cdot\gamma^{(1)}), \ldots, \exp(2\pi\sqrt{-1}\,\mathbf{n}\cdot\gamma^{(D)}))$. We first show that all these characters are distinct. Suppose $\exp(2\pi\sqrt{-1}\,\mathbf{n}\cdot\gamma^{(k)}) = \exp(2\pi\sqrt{-1}\,\mathbf{n}\cdot\gamma^{(l)})$ for all $\mathbf{n} \in \mathbb{Z}^N$. This implies that $\gamma^{(k)} \equiv \gamma^{(l)} (\mathrm{mod}\ \mathbb{Z}^N)$. Since also $\gamma^{(k)} - \gamma^{(l)} \in L \otimes \mathbb{R}$ we conclude that $\gamma^{(k)} - \gamma^{(l)} \in L$ and hence $k = l$.

Let ρ be a common point of $\mathbf{b}_I$ and $\mathbf{b}_J$ and consider the corresponding basis of local power series solutions indexed by I_ρ. With the notation from Section 6 we proceed by rewriting $\overline{g}^t H g = H$ as

$$(\overline{X}_\rho^t)^{-1}\chi^{-1}\overline{X}_\rho^t H X_\rho \chi X_\rho^{-1} = H.$$

This implies $\chi^{-1}\overline{X}_\rho^t H X_\rho \chi = \overline{X}_\rho^t H X_\rho$. Since χ consists of distinct characters on its diagonal we conclude that $\overline{X}_\rho^t H X_\rho$ is a diagonal matrix. Hence the columns of X_ρ are orthogonal with respect to H. In particular X_I, X_J are orthogonal with respect to H. □

A question is whether the orthogonality property of Proposition 7.2 suffices to determine H uniquely (up to a scalar factor). Also, to determine H completely we need to know the values of $\overline{X}_I H X_I$ for every I. So far we have not been able to answer these questions. Instead we end with the following conjecture.

Conjecture 7.3 *The orthogonality properties from Proposition 7.2 suffice to determine H up to a scalar factor. Furthermore, after suitable normalisation of H we have for every $I \in \mathcal{I}$ that $\overline{X}_I H X_I = \Delta_I \prod_{i \notin I} \sin(\pi\gamma_i^I)$.*

Consequently, this conjecture implies that the signature of the hermitian form H is determined by the signs of the values of $\prod_{i \notin I} \sin(\pi\gamma_i^I)$ as I runs through the elements of $\mathcal{I}(\rho)$ for some convergence direction ρ.

8 An example, G_3

Recall that the A-matrix for Horn's system G_3 reads

$$A = \begin{pmatrix} -1 & 0 & 1 & 2 \\ 2 & 1 & 0 & -1 \end{pmatrix}$$

and the parameter vector is $(-a, -b)^t$. The B-matrix is given by

$$B = \begin{pmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \end{pmatrix}$$

The formal solution series reads

$$\sum_{p,q} \frac{v_1^{p+\gamma_1} v_2^{-2p+q+\gamma_2} v_3^{p-2q+\gamma_3} v_4^{q+\gamma_4}}{\Gamma(p+\gamma_1+1)\Gamma(-2p+q+\gamma_2+1)\Gamma(p-2q+\gamma_3+1)\Gamma(q+\gamma_4+1)}$$

where $A(\gamma_1, \ldots, \gamma_4)^t = (-a, -b)^t$. To get a power series we choose $\gamma_1 = \gamma_4 = 0$ for example. This implies that we must take $\gamma_2 = -b, \gamma_3 = -a$. We get

$$\Phi_{0,-b,-a,0} = v_2^{-b} v_3^{-a} \sum_{p,q} \frac{(v_1 v_3 v_2^{-2})^p (v_2 v_4 v_3^{-2})^q}{\Gamma(p+1)\Gamma(-2p+q-b+1)\Gamma(p-2q-a+1)\Gamma(q+1)}.$$

This is a power series expansion in $x = v_1 v_3 / v_2^2, y = v_2 v_4 / v_3^2$ since only terms with $p, q \geq 0$ are non-zero. Using the identity $\Gamma(z)\Gamma(1-z) = \pi / \sin(\pi z)$ this implies

$$\Phi_{0,-b,-a,0} = \sin(\pi a) \sin(\pi b) v_2^{-b} v_3^{-a} \sum_{p,q} \frac{\Gamma(2p-q+b)\Gamma(-p+2q+a)}{p!q!} (-x)^p (-y)^q.$$

This is, up to a factor, the classical series $G_3(a, b, -x, -y)$.

Let us now take $\gamma_2 = \gamma_3 = 0$. Then $\gamma_1 = -(a+2b)/3$ and $\gamma_4 = -(2a+b)/3$. We get

$$\Phi_{-(a+2b)/3,0,0,-(2a+b)/3} = v_1^{-(a+2b)/3} v_4^{-(2a+b)/3} \times \sum_{p,q} c_{pq} x^p y^q$$

where the coefficients c_{pq} are product of inverse Γ-values. The c_{pq} all vanish outside the region with $-2p+q \geq 0, p-2q \geq 0$. Putting $m = -2p+q, n = p-2q$ we get that $p = -(2m+n)/3$ and $q = -(m+2n)/3$. So $x^p y^q = (x^{-2} y^{-1})^{m/3} (x^{-1} y^{-2})^{n/3}$ and we essentially have a power series in the variables $x^{-2} y^{-1}$ and $x^{-1} y^{-2}$. Similarly we have for $\epsilon \in \{-1, 1\}$,

$$\Phi_{-(a+2(b-\epsilon))/3,\epsilon,0,-(2a+(b-\epsilon))/3} = v_1^{-(a+2b-2\epsilon)/3} v_2^{\epsilon} v_4^{-(2a+b-\epsilon)/3} \times \sum_{p,q} c_{pq} x^p y^q.$$

Let us also write down all other possibilities,

$$\Phi_{0,0,-a-2b,b} = v_3^{-a-2b} v_4^{b} \times \sum_{p,q} c_{pq} x^p y^q,$$

which is a summation over all integers p, q with $-2p+q \geq, p-2q \geq 0$, and hence a power series in xy^2 and y.

$$\Phi_{a,-b-2a,0,0} = v_1^{a} v_2^{-b-2a} \times \sum_{p,q} x^p y^q,$$

which is a power series in x, x^2y. For $\epsilon \in \{0,1\}$ we have

$$\Phi_{0,-b-(a+\epsilon)/2,\epsilon,(a+\epsilon)/2} = v_2^{-b-(a+\epsilon)/2} v_3^{\epsilon} v_4^{(a+\epsilon)/2} \times \sum_{p,q} c_{pq} x^p y^q,$$

which is a summation over $p \geq, p - 2q + \epsilon \geq 0$ and essentially a power series in x, x^2y. Finally we have for $\epsilon \in \{0,1\}$,

$$\Phi_{-(b+\epsilon)/2,\epsilon,-a-(b+\epsilon)/2,0} = v_1^{-(b+\epsilon)/2} v_2^{\epsilon} v_4^{-a-(b+\epsilon)/2} \times \sum_{p,q} c_{pq} x^p y^q,$$

which is a summation over $q \geq 0, q - 2p + \epsilon \geq$ and essentially a power series in y, y^2x.

So the set $\mathcal{I}$ consists of 10 elements, 3 times $\{2,3\}$, 2 times $\{1,3\}$, 2 times $\{2,4\}$ and the three sets $\{1,4\}$, $\{1,2\}$,$\{3,4\}$. Here we numbered the columns of B by $1, 2, 3, 4$. Here is a table with the corresponding values of γ.

	J	γ^J
1	(2,3)	$(-a-2b, 0, 0, -2a-b)/3$
2	(2,3)	$(-a-2b-2, 3, 0, -2a-b-1)/3$
3	(2,3)	$(-a-2b+2, -3, 0, -2a-b+1)/3$
4	(1,3)	$(0, -b-a/2, 0, -a/2)$
5	(1,3)	$(0, -b-(a+1)/2, 1, -(a+1)/2)$
6	(2,4)	$(-b/2, 0, -a-b/2, 0)$
7	(2,4)	$(-(b+1)/2, 1, -a-(b+1)/2, 0)$
8	(1,4),	$(0, -b, -a, 0)$
9	(1,2),	$(0, 0, -a-2b, b)$
10	(3,4)	$(a, -b-2a, 0, 0)$

As noted earlier, the zonotope Z_B contains the three points $(-0.4, -0.4)$, $(-0.4, 0.6)$, $(0.6, -0.4)$ corresponding to the argument choices

$$\frac{\Theta_1}{2\pi} = (-0.4, 0, 0, -0.4),\ \frac{\Theta_2}{2\pi} = (0.6, 0, 0, -0.4),\ \frac{\Theta_3}{2\pi} = (-0.4, 0, 0, 0.6).$$

Note that we have seen to it that the second and third components are zero. We can now form the vectors X_J defined in (1), we list them in the order of the previous table and use the notation $e(x) = e^{2\pi\sqrt{-1}\ x}$.

	J	X_J
1	(2,3)	$(1, e(-(a+2b)/3), e(-(2a+b)/3))$
2	(2,3)	$(1, e(-(a+2b+2)/3), e(-(2a+b+1)/3))$
3	(2,3)	$(1, e(-(a+2b-2)/3), e(-(2a+b-1)/3))$
4	(1,3)	$(1, 1, e(-a/2))$
5	(1,3)	$(1, 1, -e(-a/2))$
6	(2,4)	$(1, e(-b/2), 1)$
7	(2,4)	$(1, -e(-b/2), 1)$
8	(1,4)	$(1, 1, 1)$
9	(1,2)	$(1, 1, e(b))$
10	(3,4)	$(1, e(a), 1)$

To write down local monodromies we select the loops c_1 and c_4, where c_i denotes the loop as we let v_i circumscribe a circle $|v_i| = r_i$ while keeping the other v_j fixed. The corresponding characters by which Φ_J changes are denoted by $\chi_{J,1}$ and $\chi_{J,4}$.

	J	$\chi_{J,1}$	$\chi_{J,4}$
1	(2,3)	$e(-(a+2b)/3)$	$e(-(2a+b)/3)$
2	(2,3)	$e(-(a+2b+2)/3)$	$e(-(2a+b+1)/3)$
3	(2,3)	$e(-(a+2b-2)/3)$	$e(-(2a+b-1)/3)$
4	(1,3)	1	$e(-a/2)$
5	(1,3)	1	$-e(-a/2)$
6	(2,4)	$e(-b/2)$	1)
7	(2,4)	$-e(-b/2)$	1
8	(1,4),	1	1
9	(1,2),	1	$e(b)$
10	(3,4)	$e(a)$	1

As an example for the action of c_1 on a local basis we take the three local basis solutions corresponding to $J = (2, 3)$. The transition matrix consists of the first three vector X_J from the above table,

$$X = \begin{pmatrix} 1 & 1 & 1 \\ e(-(a+2b)/3) & e(-(a+2b+2)/3) & e(-(a+2b-2)/3 \\ e(-(2a+b)/3) & e(-(2a+b+1)/3) & e(-(2a+b-1)/3) \end{pmatrix}$$

For the path c_1 we get the monodromy matrix

$$e(-(2a+b)/3)X \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} X^{-1} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & e(-b) \\ e(-a-b) & 0 & 0 \end{pmatrix}$$

with respect to the Mellin-Barnes basis, where $\omega = e(1/3)$. For c_4 we get

$$\begin{pmatrix} 0 & 0 & 1 \\ e(-a-b) & 0 & 0 \\ 0 & e(-a) & 0 \end{pmatrix}.$$

As a second example take the local basis corresponding to the sets $J = (1,4), (1,2), (3,4)$. The local monodromy for c_1 is given by the diagonal matrix with diagonal entries $(1,1,e(a))$ and for c_4 by $(1,e(b),1)$. The local transition matrix X reads

$$X = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & e(a) \\ 1 & e(b) & 1 \end{pmatrix}$$

For the monodromy matrices with respect to the Mellin-Barnes basis we get

$$X \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e(a) \end{pmatrix} X^{-1} = \begin{pmatrix} 0 & 1 & 0 \\ -e(a) & e(a)+1 & 0 \\ -1 & 1 & 1 \end{pmatrix}$$

and

$$X \begin{pmatrix} 1 & 0 & 0 \\ 0 & e(b) & 0 \\ 0 & 0 & 1 \end{pmatrix} X^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ -1 & 1 & 1 \\ -e(b) & 0 & 1+e(b) \end{pmatrix}.$$

In all there are four bases of local solutions yielding 8 monodromy matrices. Let G be the group generated by these matrices. By letting the other variables run through circles we essentially get elements of G again, possibly multiplied by scalars. This has to do with the homogeneity property $\Phi(\mathbf{t}^{\mathbf{a}_1}v_1, \ldots, \mathbf{t}^{\mathbf{a}_N}) = \mathbf{t}^{\alpha}\Phi(v_1, \ldots, v_N)$ for solutions of A-hypergeometric systems.

According to Proposition 7.2 there exist orthogonality relations between the vectors X_J. This can be pictured in the following graph, where we use the numbering from Table 8, and where two nodes are connected if the corresponding X_J are orthogonal according to Proposition 7.2 (it turns out that there are no more orthogonalities but this must verified).



It turns out that, up to a scalar there is a unique Hermitean matrix which makes these X_J orthogonal. Up to a scalar factor it equals

$$H = e(-(a+b)/2) \begin{pmatrix} 1+e(a+b) & -e(a+b) & -e(a+b) \\ -1 & 1+e(a+b) & -e(a) \\ -1 & -e(b) & 1+e(a+b) \end{pmatrix}.$$

9 Another example, $E(3,6)$

The system Aomoto-Gel'fand system $E(3,6)$ forms the subject of the second part of M.Yoshida's book [32]. It is the system that corresponds to configurations of six points (or lines) in $\mathbb{P}^2$. We start by giving the A-matrix of the system,

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix}$$

and the parameters $(\alpha_1, \alpha_2, 2-\alpha_4, 2-\alpha_5, 2-\alpha_6)$. The integrand of the Euler integral as defined in [7, p 607] reads

$$\frac{s^{\alpha_1-1}t^{\alpha_2-1}t_1^{1-\alpha_4}t_2^{1-\alpha_5}t_3^{1-\alpha_6}}{1-t_1(s+t+1)-t_2(s+v_1t+v_3)-t_3(s+v_2t+v_4)} ds \wedge dt \wedge dt_1 \wedge dt_2 \wedge dt_3.$$

Perform the substitutions $t_1 \to t_1/(s+t+1), t_2 \to t_2/(s+v_1t+v_3)$ and $t_3 \to t_3/(s+v_2t+v_4)$. We obtain the integrand

$$\frac{t_1^{1-\alpha_4}t_2^{1-\alpha_5}t_3^{1-\alpha_6}}{1-t_1-t_2-t_3} \prod_{i=1}^{6} (L_i)^{\alpha_i-1} ds \wedge dt \wedge dt_1 \wedge dt_2 \wedge dt_3$$

where

$$L_1 = s,\ L_2 = t,\ L_3 = 1,\ L_4 = s+t+1,$$

$$L_5 = s+v_1t+v_3,\ L_6 = s+v_2t+v_4$$

and $\alpha_3 = 3-\alpha_1-\alpha_2-\alpha_4-\alpha_5-\alpha_6$. Integration with respect to t_1, t_2, t_3 leaves us with a 2-form which is the integrand given on page 221 of [32], but with v_i instead of x^i. Thus we see that our A-matrix corresponds to a Gel'fand-Aomoto system which is associated to configurations of six lines in $\mathbb{P}^2$. The system is a four variable system of rank 6. It is irreducible if and only if none of the α_i is an integer, see [22, Prop 2]. A possible B-matrix reads

$$B = \begin{pmatrix} 1 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 1 \\ 1 & -1 & 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 1 & 0 & -1 & -1 & 0 & 1 & 0 & 0 & 0 \\ 1 & -1 & 0 & -1 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

The set $\mathcal{I}$ consists of 81 sets, and hence 81 distinct local solutions. The number of local solution bases is 108. In a straightforward manner one can check that the B-zonotope Z_B contains the points

$$p, p+(0,0,0,1), p+(1,0,0,0), p+(1,0,1,1), p+(1,1,0,1), p+(1,1,1,1)$$

where $p = (-0.9, -0.4, -0.5, -0.7)$. Hence the system $E(3,6)$ has a Mellin-Barnes basis of solutions. Computation of a set of generators for the monodromy group $lMon$ is now straightforward. Computation shows that there is a unique invariant Hermitean form (up to a scalar) for $lMon$. Presumably $lMon = Mon$. It turns out that this form satisfies Conjecture 7.3. For the computation of the signature we fix a local basis of solutions and compute the corresponding γ-parameters of the solution series. In order to choose a local basis we must specify 6 subsets of 4 column vectors $\mathbf{b}_i$ from the B-matrix whose positive cones have a point (convergence direction) in common. We choose

$$\{\mathbf{b}_5, \mathbf{b}_6, \mathbf{b}_8, \mathbf{b}_9\},\ \{\mathbf{b}_1, \mathbf{b}_5, \mathbf{b}_6, \mathbf{b}_8\},\ \{\mathbf{b}_1, \mathbf{b}_5, \mathbf{b}_7, \mathbf{b}_8\}$$

$$\{\mathbf{b}_1, \mathbf{b}_3, \mathbf{b}_5, \mathbf{b}_6\},\ \{\mathbf{b}_1, \mathbf{b}_3, \mathbf{b}_6, \mathbf{b}_7\},\ \{\mathbf{b}_1, \mathbf{b}_3, \mathbf{b}_7, \mathbf{b}_8\}$$

Each of these sets generate a positive cone which contains the point $(0.4, 0.5, 0.65, 0.7)$ in its interior. Each set represents a solution series corresponding to the following γ-parameters written in column form

$$\begin{array}{cccccc}
2-A & 0 & 0 & 0 & 0 & 0 \\
\alpha_1 & \alpha_1 & \alpha_1 & 2-\alpha_4 & 2-\alpha_4 & 2-\alpha_4 \\
\alpha_2 & 2-\alpha_1-\alpha_4 & 2-\alpha_1-\alpha_4 & 0 & 0 & 0 \\
2-\alpha_5 & 2-\alpha_5 & 3+\alpha_3 & 2-\alpha_5 & 3+\alpha_3 & 3+\alpha_3 \\
0 & 0 & 0 & 0 & -2+\alpha_1+\alpha_4 & -1-\alpha_3-\alpha_5 \\
0 & 0 & -1-\alpha_3-\alpha_5 & 0 & -2+\alpha_2+\alpha_6 & 0 \\
2-\alpha_6 & 1+\alpha_3+\alpha_5 & 0 & 1+\alpha_3+\alpha_5 & 0 & 0 \\
0 & 0 & 2-\alpha_6 & -2+\alpha_1+\alpha_4 & 0 & 2-\alpha_2-\alpha_6 \\
0 & -2+A & 0 & \alpha_2 & 2-\alpha_6 & \alpha_2
\end{array}$$

where $A = \alpha_1 + \alpha_2 + \alpha_4$ and $\alpha_3 = 3 - \alpha_1 - \alpha_2 - \alpha_4 - \alpha_5 - \alpha_6$. Let us see what Conjecture 7.3 (verified in this case) says about the signature. By contiguity we can assume that $0 < \alpha_1, \alpha_2, \alpha_4, \alpha_5, \alpha_6 < 1$. The signature is then determined by the signs of

$$\begin{array}{l}
\sin \pi A \\
\sin \pi(\alpha_1 + \alpha_4) \sin \pi(\alpha_3 + \alpha_5) \sin \pi A \\
\sin \pi(\alpha_1 + \alpha_4) \sin \pi(\alpha_3 + \alpha_5) \sin \pi \alpha_3 \\
\sin \pi(\alpha_3 + \alpha_5) \sin \pi(\alpha_1 + \alpha_4) \\
\sin \pi(\alpha_1 + \alpha_4) \sin \pi(\alpha_2 + \alpha_6) \sin \pi \alpha_3 \\
\sin \pi(\alpha_3 + \alpha_5) \sin \pi(\alpha_2 + \alpha_6) \sin \pi \alpha_3
\end{array}$$

The sum $3 - \alpha_3 = \alpha_1 + \alpha_2 + \alpha_4 + \alpha_5 + \alpha_6$ lies strictly between 0 and 5.

Let us assume it is between 0 and 1. Then all the above products are positive and we get signature $(6,0)$.

Let us assume $3 - \alpha_3$ is between 1 and 2. Then $\sin \pi \alpha_3$ is negative. Suppose $\sin \pi(\alpha_1 + \alpha_4)$ and $\sin \pi(\alpha_3 + \alpha_5)$ have opposite sign. Then the first two of the sin-products have opposite sign, as well as products number 3 and 4 and products 5 and 6. Hence the signature is $(3,3)$. Now suppose that $\sin\pi(\alpha_1+\alpha_4)$ and $\sin\pi(\alpha_3+\alpha_5)$ have the same sign. Suppose they are both positive, then necessarily, $0 < \alpha_1+\alpha_4 < 1$ and $2 < \alpha_3+\alpha_5 < 3$. From the last one we get $0 < \alpha_1+\alpha_2+\alpha_4+\alpha_6 < 1$. Hence $0 < \alpha_1+\alpha_2+\alpha_4 < 1$ and $0 < \alpha_2+\alpha_6 < 1$. Hence the signs are $1,1,-1,1,-1,-1$ so the signature is $(3,3)$. Finally suppose that $\sin\pi(\alpha_1+\alpha_4)$ and $\sin\pi(\alpha_3+\alpha_5)$ are both negative. Then, necessarily, $1 < \alpha_1+\alpha_4 < 2$ and $1 < \alpha_3+\alpha_5 < 2$. From the last one we infer $1 < \alpha_1+\alpha_2+\alpha_4+\alpha_6 < 2$. From these inequalities it follows that $1 < \alpha_1+\alpha_2+\alpha_4 < 2$ and $0 < \alpha_2+\alpha_6 < 1$. The signs now read $-1,-1,-1,1,1,1$. Again we see that the signature is $(3,3)$.

Let us assume that $3-\alpha_3$ lies between 2 and 3. Then $\sin\pi\alpha_3$ is positive. Suppose $\sin\pi(\alpha_1+\alpha_4)$ and $\sin\pi(\alpha_3+\alpha_5)$ have opposite sign. Then the signs on place 3 and 4 are -1. The first two signs will be opposite and the signs on positions 5 and 6 will also be opposite. Hence we get signature $(4,2)$. Suppose $\sin\pi(\alpha_1+\alpha_4)$ and $\sin\pi(\alpha_3+\alpha_5)$ are both positive. Then $0 < \alpha_1+\alpha_4 < 1$ and $0 < \alpha_3+\alpha_5 < 1$. The last one implies $2 < \alpha_1+\alpha_2+\alpha_4+\alpha_6 < 3$. Hence $1 < \alpha_2+\alpha_6 < 2$ and $1 < \alpha_1+\alpha_2+\alpha_4 < 2$. We get the signs $-1,-1,1,1,-1,-1$ and signature $(4,2)$. Finally suppose $\sin\pi(\alpha_1+\alpha_4)$ and $\sin\pi(\alpha_3+\alpha_5)$ are both negative. Then $1 < \alpha_1+\alpha_4 < 2$ and $1 < \alpha_3+\alpha_5 < 2$. The latter implies $1 < \alpha_1+\alpha_2+\alpha_4+\alpha_6 < 2$. And so $1 < \alpha_1+\alpha_2+\alpha_4 < 2$ and $0 < \alpha_2+\alpha_6 < 1$. Hence we get the signs $-1,-1,1,1,-1,-1$ and signature $(4,2)$.

When $3-\alpha_3$ exceeds 3 we can apply similar arguments or note that we might as well consider the system with $1-\alpha_1, 1-\alpha_2, 1-\alpha_4, 1-\alpha_5, 1-\alpha_6$ whose monodromy group is isomorphic with the complex conjugate of the original group.

Thus we conclude that, under the assumption $0 < \alpha_1,\alpha_2,\alpha_4,\alpha_5,\alpha_6 < 1$ we have the following table

$\lfloor\alpha_1+\alpha_2+\alpha_4+\alpha_5+\alpha_6\rfloor$	signature
0	$(6,0)$
1	$(3,3)$
2	$(4,2)$
3	$(3,3)$
4	$(6,0)$

This is accordance with [22, Prop 1] (except for a small printing error). Note that signature $(5,1)$ does not occur.

10 Another example, Lauricella F_D

Consider the set $A \subset \mathbb{Z}^{d+2}$ given by

$$\{\mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_{d+2}, \mathbf{e}_1+\mathbf{e}_2-\mathbf{e}_{d+2}, \mathbf{e}_1+\mathbf{e}_3-\mathbf{e}_{d+2}, \ldots, \mathbf{e}_1+\mathbf{e}_{d+1}-\mathbf{e}_{d+2}\}.$$

Here d is the actual number of classical variables, so $d = 2$ in the Appell case F_1. The parameter vector reads $(-a, -b_1, \ldots, -b_d, c-1)$.

A possible B-matrix is given by

$$B = \begin{pmatrix} -1 & -1 & 0 & \ldots & 0 & 1 & 1 & 0 & \ldots & 0 \\ -1 & 0 & -1 & \ldots & 0 & 1 & 0 & 1 & \ldots & 0 \\ \vdots & & & & & & & & & \vdots \\ -1 & 0 & 0 & \ldots & -1 & 1 & 0 & 0 & \ldots & 1 \end{pmatrix}$$

Denote the columns by $\mathbf{b}_1, \mathbf{b}_2, \ldots, \mathbf{b}_{2d+2}$. Since the last $d+1$ columns equal minus the first $d+1$ columns, the B-zonotope consists of all points

$$\{\lambda_1 \mathbf{b}_1 + \lambda_2 \mathbf{b}_2 + \cdots + \lambda_{d+1} \mathbf{b}_{d+1} | -1/2 < \lambda_j < 1/2\}$$

Consider the point $(x_1, \ldots, x_d)$ in Z_B with $1/2 > x_1 > x_2 > \cdots > x_d > 0$. Then the d points $(x_1 - 1, x_2, \ldots, x_d)$, $(x_1 - 1, x_2 - 1, x_3, \ldots, x_d)$ untill $(x_1 - 1, x_2 - 1, \ldots, x_d - 1)$ are also in Z_B. Hence there exists a Mellin-Barnes basis of solutions for the Lauricella system F_D.

We now select a local basis of solutions. Choose a convergence direction $\rho = (\rho_1, \ldots, \rho_d)$ with $0 < \rho_1 < \rho_2 < \ldots < \rho_d$ in the column space of B. Then

$$\begin{aligned} (\rho_1, \ldots, \rho_d) &= \rho_1 \mathbf{b}_{d+3} + \rho_2 \mathbf{b}_{d+4} + \cdots + \rho_d \mathbf{b}_{2d+2} \\ &= \rho_1 \mathbf{b}_{d+2} + (\rho_2 - \rho_1) \mathbf{b}_{d+4} + \cdots + (\rho_d - \rho_1) \mathbf{b}_{2d+2} \\ &= (\rho_2 - \rho_1) \mathbf{b}_2 + \rho_2 \mathbf{b}_{d+2} + (\rho_3 - \rho_2) \mathbf{b}_{d+5} + \cdots + (\rho_d - \rho_2) \mathbf{b}_{2d+2} \\ &\vdots \\ &= (\rho_d - \rho_1) \mathbf{b}_2 + (\rho_d - \rho_2) \mathbf{b}_3 + \cdots + (\rho_d - \rho_{d-1}) \mathbf{b}_d + \rho_d \mathbf{b}_{d+2} \end{aligned}$$

All coordinates are positive and so the set $\mathcal{I}(\rho)$ consists of $d+1$ sets of indices, written as rows in the following matrix,

$$\begin{array}{ccccc} d+3 & d+4 & d+5 & \ldots & 2d+2 \\ d+2 & d+4 & d+5 & \ldots & 2d+2 \\ d+2 & 2 & d+5 & \ldots & 2d+2 \\ d+2 & 2 & 3 & \ldots & 2d+2 \\ \vdots & & & & \vdots \\ d+2 & 2 & 3 & \ldots & d \end{array}$$

The corresponding parameter vectors γ read as follows,

$$\begin{array}{cccccccccc} -a & -b_1 & -b_2 & \ldots & -b_d & c' & 0 & 0 & \ldots & 0 \\ c'-a & c'-b_1 & -b_2 & \ldots & -b_d & 0 & -c' & 0 & \ldots & 0 \\ c'-a & 0 & c'-b_1-b_2 & \ldots & -b_d & 0 & -b_1 & b_1-c' & \ldots & 0 \\ \vdots & & & & & & & & & \vdots \\ c'-a & 0 & 0 & \ldots & c' - \sum_{j=1}^{d} b_j & 0 & -b_1 & -b_2 & \ldots & -c' + \sum_{j=1}^{d-1} b_j \end{array}$$

where $c' = c - 1$. Conjecture 7.3 now implies that the signs of the products

$$\begin{aligned}
&\sin\pi(-a)\sin\pi(-c)/\sin\pi(a-c)\\
&\sin\pi(-c)\sin\pi(b_1-c)/\sin\pi b_1\\
&\sin\pi(b_1-c)\sin\pi(b_1+b_2-c)/\sin\pi b_2\\
&\vdots\\
&\sin\pi(b_1+\cdots+b_{d-1}-c)\sin\pi(b_1+\cdots+b_d-c)/\sin\pi b_d
\end{aligned}$$

determine the signature of the invariant Hermitean form. Let us replace c by $a-b_0$. Suppose in addition that $0 < b_j < 1$ for $j = 0, 1, \ldots, d$ (by contiguity). Then the signature of the hermitian form is determined by the signs of the numbers

$$\sin\pi\left(-a+\sum_{j=0}^{k} b_j\right)\sin\pi\left(-a+\sum_{j=0}^{k-1} b_j\right)$$

for $k = 0, 1, 2, \ldots, d$. These numbers are negative if there is an integer q such that $-a+\sum_{j=0}^{k-1} b_j < q < -a+\sum_{j=0}^{k} b_j$. Hence the number of negative signs equals $\lfloor -a+\sum_{j=0}^{d} b_j\rfloor - \lfloor -a\rfloor$. Assuming $0 < -a < 1$, as we may, we see that the signature equals $d+1-\mu, \mu$ where $\mu = \lfloor -a+b_0+\cdots+b_d\rfloor$.

In the work of Picard [25], Terada [31] and Deligne-Mostow [9] one assumes the additional existence of $0 < b_{d+2} < 1$ (the exponent at ∞ of the multivalued differential form) such that $\sum_{i=0}^{d+2} b_i = 2$. In that case it follows from our consideration that the signature equals $(1, d)$, as is the case in the papers just mentioned.

References

[1] A.Adolphson, Hypergeometric functions and rings generated by monomials. Duke Math. J. **73** (1994), 269-290.

[2] G.E.Andrews,R.Askey,R.Roy, *Special Functions*, Encyclopedia of Math and its applications 71, Cambridge, 1999.

[3] I.A.Antipova, Inversion of multidimenional Mellin transforms, Communications of the Moscow Math Society, Uspekhi Mat. Nauk **62** (2007), 147-148; Russian Math. Surveys **62** (2007), 977-979.

[4] I.A.Antipova, Inversion of many-dimensional Mellin transforms and solutions of algebraic equations, Math Sbornik **198** (2007), 474-463.

[5] F.Beukers, *Notes on A-hypergeometric functions*, to appear in Société mathématique de France, Séminaires et Congrès.

[6] F.Beukers, Irreducibility of A-hypergeometric systems, (2010) arXiv:1007.4644[math.AG], to appear in Indag.Math.

[7] F.Beukers, Algebraic A-hypergeometric functions, Invent. Math. **180** (2010), 589-610.

[8] F.Beukers, G.Heckman, Monodromy for the hypergeometric function ${}_nF_{n-1}$. Invent. Math. **95** (1989), 325-354

[9] P.Deligne, G.D.Mostow, Monodromy of hypergeometric functions and non-lattice integral monodromy, Publ. Math. IHES **63** 1986.

[10] I.M.Gelfand, M.I.Graev, A.V.Zelevinsky, Holonomic systems of equations and series of hypergeometric type, Doklady Akad. Nauk SSSR **295** (1987), 14-19 (in Russian).

[11] I.M.Gelfand, A.V.Zelevinsky, M.M.Kapranov, Equations of hypergeometric type and Newton polytopes, Doklady Akad. Nauk SSSR **300** (1988), 529-534 (in Russian)

[12] I.M.Gelfand, A.V.Zelevinsky, M.M.Kapranov, Hypergeometric functions and toral manifolds, Functional Analysis and its applications **23** (1989), 94-106.

[13] I.M.Gelfand, M.M.Kapranov, A.V.Zelevinsky, Generalized Euler integrals and A-hypergeometric functions, Adv. in Math **84** (1990), 255-271.

[14] I.M.Gelfand, M.M.Kapranov, A.V.Zelevinsky, A correction to the paper "Hypergeometric equations and toral manifolds". Functional Analysis and its applications, **27** (1993), p295.

[15] V. Golyshev, A. Mellit, Gamma structures and Gauss's contiguity, (2009) arXiv:0902.2003v1 [math.AG]

[16] Y.Haraoka, Y.Ueno, Rigidity for Appell's hypergeometric series F_4, Funkcialaj Ekvasioj **51** (2008), 149-164.

[17] J.Kaneko, Monodromy group of Appell's system F_4, Tokyo J. Math **4** (1981), 35-54.

[18] M.Kato, Appell's hypergeometric systems F_2 with finite irreducible monodromy groups. Kyushu J. Math. **54** (2000), 279-305.

[19] M.Kita, M.Yoshida, Intersection theory for twisted cycles, Math.Nachr. **166** (1994), 287-304, part II in Math.Nachr. **168** (1994), 171-190.

[20] N.W.Maclachlan, *Complex variable theory and transform calculus*, second edition, Cambridge Univ. Press 1953.

[21] K.Matsumoto, T.Sasaki, N.Takayama, M.Yoshida, Monodromy of the hypergeometric equation of type $(3, 6)$, I, Duke Math.J. **71** (1993), 403-426.

[22] K.Matsumoto, T.Sasaki, N.Takayama, M.Yoshida, Monodromy of the hypergeometric equation of type $(3,6)$, II, The unitary reflection group of order $2^9 \cdot 3^7 \cdot 5 \cdot 7$, Ann. Sc. Norm. Sup. Pisa **20** (1993), 617-631.

[23] K.Mimachi, Intersection numbers for twisted cycles and the connection problem associated with the generalized hypergeometric function ${}_{n+1}F_n$, International Math. Research Notes 2010.

[24] L.Nilsson, *Amoebas, Discriminants, and Hypergeometric Functions*, PhD dissertion Stockholm University, 2009.

[25] E.Picard, Sur une extension aux fonctions de deux variables du problème de Riemann relatif aux fonctions hypergéométriques. Ann. Ec. Norm. Sup. II, **10** (1881), 304-322.

[26] M.Saito, B.Sturmfels, N.Takayama, *Gröbner Deformations of Hypergeometric Differential Equations*, Algorithms and Computation in Math. 6, Springer 2000

[27] T.Sasaki, On the finiteness of the monodromy group of the system of hypergeometric differential equations (F_D). J. Fac. Sci. Univ. Tokyo Sect. IA Math. **24** (1977) 565-573.

[28] F.C.Smith, Relations among the fundamental solutions of the generalized hypergeometric equation when $p = q+1$. Non-logarithmic cases. Bulletin of the AMS **44** (1938), 429-33.

[29] J.Stienstra, GKZ hypergeometric structures, in: *Arithmetic and geometry around hypergeometric functions*, 313-371, Progr. Math., 260, Birkhäuser, Basel, 2007.

[30] K.Takano, Monodromy of the system for Appell's F_4, Funkcialaj Ekvasioj **23** (1980), 97-122.

[31] T. Terada; Fonctions hypergéométriques F1 et fonctions automorphes I. J. Math. Soc. Japan **35** (1983), 451-475.

[32] M.Yoshida, *Hypergeometric Functions, my Love*, Aspects of Mathematics **32**, Vieweg Verlag 1997.

[33] O.N.Zhdanov, A.K.Tsikh, Studying the multiple Mellin-Barnes integral by means of multidimensional residues, Siberian J.Math **39** (1998), 245-260.